

12 LEVEL III

AD A072174

DDC  
RECEIVED  
AUG 3 1979  
B

Disclaimer

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Disposition

Destroy this report when it is no longer needed. Do not return it to the originator.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM												
1. REPORT NUMBER ARCSL-TR-79019	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER												
4. TITLE (and Subtitle) IONIC CLUSTER MASS SPECTROMETRY. 1. DETECTION AND IDENTIFICATION OF AIRBORNE ORGANIC VAPORS AND ESTABLISHMENT OF OPERATIONAL MECHANISMS IN THE IONIZATION DETECTOR SYSTEM		5. TYPE OF REPORT & PERIOD COVERED Technical Report June 1972-April 1974												
7. AUTHOR(s)  Charles S. Harden Thomas C. Imeson		6. PERFORMING ORG. REPORT NUMBER												
9. PERFORMING ORGANIZATION NAME AND ADDRESS Commander/Director, Chemical Systems Laboratory ✓ Attn: DRDAR-CLB-CA Aberdeen Proving Ground, Maryland 21010		8. CONTRACT OR GRANT NUMBER(s)												
11. CONTROLLING OFFICE NAME AND ADDRESS Commander/Director, Chemical Systems Laboratory Attn: DRDAR-CLJ-R Aberdeen Proving Ground, Maryland 21010		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS  Task 1W762710AD2901												
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE June 1979												
		13. NUMBER OF PAGES 25												
		15. SECURITY CLASS. (of this report)  UNCLASSIFIED												
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE NA												
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.														
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)  DDC RECEIVED AUG 3 1979 B														
18. SUPPLEMENTARY NOTES														
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  <table border="0"> <tr> <td>Mass spectrometry</td> <td>Chemical ionization</td> <td>Dimethyl sulfoxide</td> </tr> <tr> <td>Ionic clusters</td> <td>Atmospheric pressure ionization</td> <td>Ionization detector</td> </tr> <tr> <td>Ion-molecule reactions</td> <td>Diisopropyl methylphosphonate</td> <td>Diffusion</td> </tr> <tr> <td>Detection</td> <td>Dimethyl methylphosphonate</td> <td></td> </tr> </table>			Mass spectrometry	Chemical ionization	Dimethyl sulfoxide	Ionic clusters	Atmospheric pressure ionization	Ionization detector	Ion-molecule reactions	Diisopropyl methylphosphonate	Diffusion	Detection	Dimethyl methylphosphonate	
Mass spectrometry	Chemical ionization	Dimethyl sulfoxide												
Ionic clusters	Atmospheric pressure ionization	Ionization detector												
Ion-molecule reactions	Diisopropyl methylphosphonate	Diffusion												
Detection	Dimethyl methylphosphonate													
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  <p>An analytical technique called Ionic Cluster Mass Spectrometry (ICMS) has been developed. The technique is akin to chemical ionization mass spectrometry in that both concepts rely upon the mass analysis of ions that are products of ion-molecule reactions occurring in a reactant gas in the ion source. ICMS employs moist air as the reactant gas at ion source pressures of 10 mmHg to 1 atmosphere. Samples can be injected directly into the ion source in a clean, moist airstream or ambient atmosphere may constitute the reactant gas plus sample.</p> <p>(Continued on reverse side)</p>														

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

20. ABSTRACT (Continued)

*A* ICMS analyses of organic vapors in air have been used to aid in the establishment of response mechanisms in the Ionization Detector System (IDS). Ionic clusters to which the IDS respond have been identified. The conclusion drawn from these studies is that differential diffusion of ionic clusters from an airstream predominates in the physics of the response mechanism.

*A*

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

## PREFACE

This work was authorized under Project/Task 1W762710AD2901, Chemical Detection and Identification Technology, Detection and Identification Concepts. The work covered by this report began in June 1972 and terminated in April 1974.

Reproduction of this document in whole or in part is prohibited except with permission of the Commander/Director, Chemical Systems Laboratory, Attn: DRDAR-CLJ-R, Aberdeen Proving Ground, Maryland 21010. However, Defense Documentation Center and the National Technical Information Service are authorized to reproduce the document for US Government purposes.

## Acknowledgment

The authors acknowledge the technical efforts of J. A. Parsons and D. J. Buckley who gathered most of the experimental data.

Accession for	
NTIS GARY	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Chemical	<input type="checkbox"/>
June 1974	<input type="checkbox"/>
By	
Distribution	
Availability Code	
Dist	Avail. for special
A	

## CONTENTS

	<u>Page</u>
I. INTRODUCTION . . . . .	7
II. EXPERIMENTAL . . . . .	8
III. RESULTS AND DISCUSSION . . . . .	10
A. Ionic Mass Spectrometry . . . . .	10
B. Ionization Detector System . . . . .	14
1. Qualitative Response Studies . . . . .	14
2. Diffusion Studies . . . . .	16
IV. SUMMARY . . . . .	20
A. Ionic Cluster Mass Spectrometry . . . . .	20
B. Ionization Detector System . . . . .	20
LITERATURE CITED . . . . .	21
DISTRIBUTION LIST . . . . .	23

## LIST OF FIGURES

### Figure

1	Schematic Diagram of the ICMS Apparatus . . . . .	8
2	Schematic Diagram of an IDS Cell . . . . .	9
3	Fractional Ion Intensities in DMMP Versus Pressure . . . . .	11
4	Fractional Ion Intensities in DMSO Versus Pressure . . . . .	11
5	ICMS Relative Ion Intensities Versus [DMMP] . . . . .	13
6	IDS Response Versus Concentration at Several Flow Rates . . . . .	15
7	Positive and Negative IDS Response As a Function of Flow Rate . . . . .	19

IONIC CLUSTER MASS SPECTROMETRY. I. DETECTION AND  
IDENTIFICATION OF AIRBORNE ORGANIC VAPORS AND  
ESTABLISHMENT OF OPERATIONAL MECHANISMS IN  
THE IONIZATION DETECTOR SYSTEM

I. INTRODUCTION.

The study of proton ion/dipole clusters in the gas phase<sup>1-4</sup> has been extended in connection with the search for new concepts in chemical agent (and other atmospheric contaminants) detection. The result has been an analytical technique called Ionic Cluster Mass Spectrometry (ICMS).<sup>5</sup> ICMS is akin to the technique of chemical ionization (CI) mass spectrometry in that both concepts rely upon the mass analysis of ions that are products of ion-molecule reactions occurring in a reactant gas in the ion source. At atmospheric pressure in the ion source the technique has been termed Atmospheric Pressure Ionization (API) Mass Spectrometry.<sup>6</sup> Users of API must take care to insure that the ion source is as free as possible of water vapor which will interfere with the ionization of the ultratrace amounts of samples to be analyzed. ICMS employs moist air as the reactant gas at ion source pressures of 10 mmHg to 1 atmosphere. The sample may be introduced into a clean, moist airstream or the sample plus reactant gas may constitute ambient atmosphere. Sample/reactant gas ratios may be as low as 5 mole parts in  $10^{10}$ .

During the study to date of ICMS no fragmentation of sample molecules has been observed, only increases in sample mass by protonation and hydration. Since only cluster ions are observed the term ICMS is more descriptive of this mass-spectrometric-analysis technique.

Mechanisms, rate constants, and enthalpies of formation for ion clusters  $H^+(H_2O)_n$  (derived from ion molecule reactions initiated by the primary ions  $N_2^+$  and  $O_2^+$ ) have been established where the primary ionization source has been electron irradiation,<sup>7,8</sup> in flowing afterglow experiments,<sup>9</sup> and in corona discharges.<sup>5,10,11</sup> For the ICMS technique, the hydrated proton species serve as precursors for clusters  $H^+(H_2O)_n(A)_m$  which represent the qualitative and quantitative indicators of the trace compound A. The mechanism of formation of  $H^+(H_2O)_n(A)_m$  is not dependent upon the method of formation of primary ions, i.e. electron impact (EI) or electrical discharges. The sensitivity of the technique is, however, dependent upon the rate of formation of the primary ions, the water vapor concentration, and the pressure of the reactant gas. A corona discharge was selected for most of this work because it is an extremely intense source of primary ions. A tritium  $\beta$  source proved to be a much more stable source of ions.

Some of the results of the ICMS studies have been applied to the establishment of the response mechanisms in the Ionization Detector System (IDS) now under development for the Army and the Air Force. The basic field unit, as it exists to date, has been developed by Minneapolis Honeywell, Inc. The mechanism by which the IDS selectively responds to chemical agents in the atmosphere has not been fully understood. The construction details, sensitivity, and specificity of the system have been given in a contractor report.<sup>12</sup>

## II. EXPERIMENTAL.

Figure 1 is a diagram of the ICMS apparatus. Ions produced in a corona discharge or as a result of ion-molecule reactions (formed at room temperature,  $\sim 25^{\circ}\text{C}$ ) diffuse through a pinhole into a vacuum chamber where they are focused into a quadrupole mass spectrometer. Ion detection is accomplished with an electron multiplier coupled to a high-speed picoammeter or electrometer amplifier.




Figure 1. Schematic Diagram of the ICMS Apparatus

Air or nitrogen containing trace quantities of water vapor and the substance to be analyzed flow into the ion source -- a corona discharge cylinder at C. The discharge is established between a 90% Pt - 10% Rh wire, 0.08-mm-diameter, and a stainless steel cylinder, 12-mm-ID, located coaxial to one another. The discharge current is maintained at  $10\ \mu\text{A}$  (for all source pressures) with a 6-kv variable power supply placed in series with a 4.7-megohm current-limiting resistor. Ions exit the source chamber by diffusion through an orifice, 35 microns in diameter, located in the center of S and are subsequently mass-analyzed.

Flow rate through the ion source is held constant for all pressures. Source pressures are varied to allow a control of the collision rate and the subsequent appearance and/or



disappearance of ionic species of interest. Analytical data is obtained when the source pressure is held constant and the sample concentration is variable.

A sketch of a detection cell in the IDS in the configuration used in this work is shown in figure 2. The system operates at atmospheric pressure. Air, heated to 60°C (the normal operating temperature of the IDS), is pumped through the cell at any desired flow rate. The cell is placed in a small oven so that 60°C is maintained throughout the cell. The airstream passes over a 1-Curie  $\beta$ -source (titanium tritide coated foil). The primary ions,  $N_2^+$ ,  $O_2^+$ , and  $O_3^+$ , are formed near the  $\beta$ -source by electron impact or electron attachment. A sequence of ion-molecule reactions follows and equilibrium among ionic clusters is rapidly established. The mixture of air and ions then flows through a series of baffles to a Faraday cup ion collector. One end of the Faraday cup is a grid to allow the air to flow through. The  $\beta$ -source which is in electrical contact with the center manifold stud can be biased either positively, negatively, or maintained at zero potential with respect to the collector. The ion currents are measured with a picoammeter.




Figure 2. Schematic Diagram of an IDS Cell

Trace concentrations of samples are prepared in a continuous flow system by using air dilution of vapor from the compound of interest.

### III. RESULTS AND DISCUSSION.

#### A. Ionic Mass Spectrometry.

Specific ion species indicated in an ionic cluster mass spectrum of dimethyl methylphosphonate are given in the table below.

Identification of DMMP Spectral Elements

<u>Ion</u>	<u>m/e</u>	<u>I/ΣI (%)</u>
$H^+(H_2O)_3$	55	2.5
$H^+(H_2O)_4$	73	8.0
$H^+(H_2O)_5$	91	7.0
$H^+(H_2O)_6$	109	1.5
$H^+(DMMP)$	125	11.1
$H^+(H_2O)_7$	127	1.0
$H^+(DMMP)(H_2O)$	143	14.1
$H^+(DMMP)(H_2O)_2$	161	8.5
$H^+(DMMP)_2$	249	43.2
Other	-	3.1

This spectrum was taken using air as the reactant gas at a source pressure of 140 mmHg,  $[DMMP] = 0.6$  ppm, and  $[H_2O] \sim 100$  ppm. As can be seen in the table, indicators of the trace compound appear at m/e of  $[MW + 1 + n(18)]$ , e.g.,  $H^+(DMMP)(H_2O)_n$ , and at m/e of  $[n(MW) + 1]$ , e.g.,  $H^+(DMMP)_n$ , where n is 1 or 2. A cluster containing a single organic molecule is referred to as a monomer or hydrated monomer if it contains water; those clusters containing two organic molecules are called dimers.

Ions that appear under the category "other" in the table constitute background and can be attributed to clusters formed with methanol and acetone - solvents used in cleaning the apparatus. Consequently, no fragment ions need or can be assigned to the spectral elements.

Figures 3 and 4 illustrate the normalized ion intensities versus pressure of DMMP and dimethyl sulfoxide (DMSO) systems in the pertinent high-pressure regions. Concentrations of DMMP and DMSO are approximately the same (0.6 and 0.7 ppm respectively) and  $[H_2O] \sim 100$  ppm. It will be noted that indicators of DMMP appear at lower pressures than those for

DMSO and that the rates of formation of  $\text{H}^+(\text{DMSO})_2$  and  $\text{H}^+(\text{DMSO})$  are about the same. Without evidence to the contrary, the mechanism of formation of ion clusters for both DMMP and DMSO is assumed to be the same. However, for ion clusters formed from DMMP, the formation of the proton-dimer has predominance over the formation of the proton-monomer, while for DMSO these two reactions have a parity in their probability of occurrence.




Figure 3. Fractional Ion Intensities in DMMP Versus Pressure





Figure 4. Fractional Ion Intensities in DMSO Versus Pressure

Dipole moments for DMMP<sup>13</sup> and DMSO<sup>14</sup> are respectively 3.62 and 3.96 Debye units; whereas their polarizabilities, computed from refractive indices, are 10.6 Å<sup>3</sup> and 7.9 Å<sup>3</sup>. An explanation for the apparent greater stability and ease of formation of clusters containing DMMP compared to DMSO may be interpreted as a greater influence of polarizability over dipole moment. However, the refractive indices and densities of the two substances are such that the difference in their polarizabilities is primarily a function of their molecular weights. Consequently, while dipole moment and polarizability certainly influence cluster formation, it is assumed that the greater number of vibrational degrees of freedom resident in the DMMP molecule impart a larger stability to its clusters when compared to those of DMSO. This indicates, that within certain unspecified limits, the larger a dipolar molecule the more stable and readily formed will be its ion clusters.


The quantification of the ICMS technique and its sensitivity with respect to trace atmospheric impurities is determined by studying ion cluster formation as a function of the impurity concentration at constant pressure. Figure 5 illustrates the dependence of various ionic cluster concentrations upon DMMP at 140 mmHg. The plotted points indicate experimental data and the smooth curves represent the results of numerically fitting kinetic equations to the data. For the curve-fitting procedure, several reaction mechanisms were considered. The mechanism resulting in the best fit is slightly different from one previously proposed<sup>5</sup> but, nevertheless, is consistent with the qualitative predictions that can be drawn from figures 3 and 4. This mechanism and calculated rate constants are given below.



$$k = 1.85 \times 10^{-8} \text{ cm}^3 \text{ sec}^{-1}$$



$$k = 4.06 \times 10^{-8} \text{ cm}^3 \text{ sec}^{-1}$$



$$k_f = 2.77 \times 10^{-13} \text{ cm}^3 \text{ sec}^{-1}$$

$$k_r = 7.4 \times 10^{-25} \text{ cm}^3 \text{ sec}^{-1}$$

The average deviation of the calculated points from experimental data is 12.4%.




Figure 5. ICMS Relative Ion Intensities Versus [DMMP]

At concentrations near the threshold sensitivity the indicators of the trace impurities are  $H^+(A)(H_2O)$ , where A is an impurity molecule. At higher concentrations, a few ppm, the ionic cluster mass spectrum consists only of the ionic clusters  $H^+(A)_n$ , where n is usually 1 and 2. The best sensitivity that has been observed, to date, is 0.5 ppb of diisopropyl methylphosphonate (DIMP).

#### B. Ionization Detector System.

It is now well understood that the charge carriers in ionized air, at or near atmospheric pressure, are in the form of ionic clusters, regardless of the source of primary ionization. It is, therefore, evident that responses in the IDS are caused by changes in densities of ionic clusters as a function of trace impurity concentration as an flows through the detection cell (figure 2). The current changes usually appear as increases (increase in positive ion cluster concentration) when an impurity is introduced into a clean moist airstream.

Both positive and negative primary ions are formed in the  $\beta$ -source region of the detector cell and react with neutral molecules to form positively and negatively charged clusters. In the absence of atmospheric impurities the positive clusters are  $H^+(H_2O)_n$  and the negative clusters are primarily  $O_2^-(H_2O)_n$ .<sup>15</sup> Under these conditions, more positive clusters than negative clusters reach the collector per unit time. The net result is a positive current measured at the collector.

When impurities are present in the air, reactions between the impurity molecules and the hydrate clusters result in the replacement of water in the clusters by the impurity. The clusters thus produced are larger in size and, presumably, more stable.

#### 1. Qualitative Response Studies.

At the beginning of a series of parametric studies on the IDS, qualitative "sniff" tests were run for approximately 60 compounds. In general, the IDS responses were positive, i.e., there were increases in the net positive cluster concentrations at the collector. Some compounds showed an increase in the net negative ion current (examples are most chloro compounds) and some showed no change in cell current.

Examination of the responses of an homologous series of primary alcohols (methyl through n-butyl) showed that the larger the impurity molecule, the larger the cell response. It was also noted that large concentrations of ammonia caused no cell response. Ammonia is known to react very well to form ammonia-water ion clusters<sup>14</sup> and the cluster size distribution is essentially the same as that with only water present. The conclusions drawn from these observations are (a) the larger molecules form larger clusters and (b) since larger clusters have small diffusion velocities normal to the flow streamlines they have a greater probability of surviving the flow region and increasing the collector current.

The conclusions derived from the qualitative observations were tested by measuring the response of the IDS to quantitative mixtures of several homologous series of impurity compounds.

Primary alcohols (methanol through n-pentanol) were studied at concentrations of 25 and 50 ppm. At each concentration the order of response was n-pentanol > n-butanol > n-propanol > ethanol > methanol. An ICMS analysis of methanol, ethanol, and n-propanol, each at a concentration of 25 ppm in air, shows that the alcohols readily react to form ionic clusters  $H^+(ROH)_n(H_2O)_m$  and that the average mass of the clusters increases with increasing molecular weight.

A series of CW agent simulants (phosphonate esters) was studied as functions of concentration and flow rate. The series consisted of diisopropyl methylphosphonate (MW 180), diethyl ethylphosphonate (MW 166), dimethyl methylphosphonate (MW 124), and dimethyl hydrogenphosphonate (MW 110). The results for DIMP and DMMP are shown in figure 6. The other members of the series were left out for the sake of clarity. As can be seen, the response increases to saturation with concentration. At any given concentration and flow rate the response due to DIMP > DEEP > DMMP > DMHP.




Figure 6. IDS Response Versus Concentration at Several Flow Rates

Three amines were tested quantitatively and the same behavior was noted; namely, the response of diisopropyl amine > diethyl amine >> ammonia.

The IDS response shows different response characteristics to acids. At low concentrations the response decreases (increases negatively) markedly. As concentration is increased further the response passes through a minimum and then increases toward saturation. The negative ion clusters are formed faster at lower concentrations and tend toward saturation sooner than the positive ions. The magnitude of responses (both positive and negative) due to the acids studied is butyric acid > acetic acid.

All of the quantitative work done so far shows that, within a homologous series, the larger molecules result in larger ion clusters and procedure greater IDS responses. Thus, the quantitative studies seem to verify that the size of the ionic clusters, therefore, diffusion, plays a predominant role in the IDS response mechanism.

## 2. Diffusion Studies.

The effect of cluster size can be demonstrated by studying the IDS response to water vapor at various airflow rates. The positive  $I^+$ , negative  $I^-$ , ion currents (+ and - 200 volts source bias) were measured as a function of flow rate,  $Q$ , up to about 8.5 l/min. The relationship between  $I$  and  $Q$  was empirically found to be

$$I = K \cdot Q^{3/2}$$

In the vicinity of the source of ionization the concentrations of positive and negative ions is dependent upon temperature, pressure, and  $\beta$ -source activity. These parameters are held constant so that the ion density in the ion source region is constant. The ion loss mechanism in this region of the detector is primarily recombination which reduces both positive and negative ion concentrations by the same amount.

To a first approximation the cell geometry is considered as a straight cylindrical tube of volume  $V$  (the effective internal volume of the baffle section of the cell) and radius  $r$  (half the average spacing of the various components of the baffle section). A certain number of ions per unit area,  $N_0$  is introduced into the baffle section. Only radial diffusion is considered since axial diffusion does not result in a loss of ions. The one dimensional diffusion equation with the approximation for small distance is

$$n(x, t) = \frac{N_0}{2\sqrt{\pi Dt}} \exp\left[-\frac{x^2}{4Dt}\right] \approx \frac{N_0}{2\sqrt{\pi Dt}}$$

where

$n(x, t)$  = the ion concentration at a radial distance  $x$  from the tube axis at time  $t$  after the initial introduction of ions;

$D$  = the diffusion coefficient.



An ion is considered to be lost and therefore not able to reach the ion collector when it has diffused through the distance,  $r$  (the radius of the approximating tube). The ion concentration at the collector end of the tube is then given by

$$n(t) = \frac{N_0}{2\sqrt{\pi Dt}}$$

Integration of the above along the radial distance of diffusion provides the number of ions per unit area at the collector.

$$\int_0^r n(t) dx = N(t) = \int_0^r \frac{N_0 dx}{2\sqrt{\pi Dt}} = \frac{N_0 r}{2\sqrt{\pi Dt}}$$

The equality of ratios for ions per unit area,  $N$ , and ion concentration,  $n$ , can be made

$$\frac{N(t)}{N_0} = \frac{n(t)}{n_0} = \frac{r}{2\sqrt{\pi Dt}}$$

Therefore,

$$n(t) = \frac{n_0 r}{2\sqrt{\pi Dt}}$$

The ion current at the collector is

$$I = n(t) q Q$$

where

$q$  = the electronic charge ( $1.6 \times 10^{-19}$  coulomb).

Therefore,

$$I = \frac{n_0 r q Q}{2\sqrt{\pi Dt}}$$

the time,  $t$ , is the residence time of particles in the tube and is  $V/Q$ . Substituting for  $t$ , the positive or negative ion current at the collector is

$$I = \frac{n_0 r q Q^{3/2}}{2\sqrt{\pi D V}}$$

Thus, the average ionic diffusivity can be determined from the slope,  $K$  of a plot of  $I$  versus  $Q^{3/2}$ . In a typical IDS cell,  $r = 0.038$  cm and  $V = 0.19$  cm<sup>3</sup>. Dimensions of the ion source region are such that recombination of positive and negative ions is an efficient mechanism for ion loss when ion concentrations are greater than  $1.5 \times 10^7$  ion/cm<sup>3</sup>; below this concentration recombination is unimportant. Therefore, the value chosen for  $n_0$  is  $1.5 \times 10^7$  ions/cm<sup>3</sup> for both positive and negative ions.

Then,

$$D = \left[ \frac{n_0 r q}{2K} \right]^2 \cdot \frac{1}{\pi V} = \frac{1.61 \cdot 10^{-23}}{K^2}$$

where

$I$  is measured in amperes and  $Q$  in l/min.

Figure 7 represents the results of diffusion studies ( $I$  versus  $Q^{3/2}$ ) for moist air. At low humidity, separate and distinct curves are obtained for positive and negative ions. This indicates that the negative ions are smaller than the positive ions. When the humidity is increased, ionic diffusivity decreases (larger ions are formed), and the curves coincide; oppositely charged ions are essentially the same size. The values of diffusivities calculated from these data agree very well, considering the complexity of the actual cell geometry, with typical ion diffusivities ( $0.06$  cm<sup>2</sup>/sec) reported in the literature,<sup>16</sup> and are summarized below.

<u>Relative humidity (25°C)</u>	<u><math>D^+</math> (cm<sup>2</sup>/sec)</u>	<u><math>D^-</math> (cm<sup>2</sup>/sec)</u>
15%	0.075	0.094
63%	0.052	0.052
92%	0.030	0.030

The observed decrease in  $D$  with humidity is consistent with the well-known fact that the average size of hydrated-ion clusters increases with  $[H_2O]$ .




Figure 7. Positive and Negative IDS Response As a Function of Flow Rate

The conclusions drawn from these studies is that differential diffusion of ionic clusters from an airstream predominates in the physics of the response mechanism. An independent study of the IDS mechanism\* has substantiated the conclusions of this work with respect to the diffusion mechanism.

\*McClure, B. T., and Waletzko, J. A. Ionization Detector System (IDS) Mechanism. Unpublished work. Honeywell, Inc., Minneapolis, Minnesota.

#### IV. SUMMARY.

##### A. Ionic Cluster Mass Spectrometry.

Irrespective of the subtle aspects associated with ion cluster stability, the fact remains that the techniques producing them offers an opportunity for application to the detection and analysis of trace compounds in the gas phase. The ability to use atmospheric gases as reactants considerably extends the practical applicability of this technique. Its incorporation with gas chromatographic analysis of trace compounds (where  $N_2$  is a common carrier gas) has many possibilities. Several features that enhance its general attractiveness are: (1) the absence of a spectrum complicated with fragment ions, (2) high mass peaks directly related to the molecular weight of the compounds, and (3) sensitivity well below the ppm concentration range.

##### B. Ionization Detector System.

It has been shown that the charge carriers in the IDS are ionic clusters of water and trace gas molecules. The ability to use such a system as a detector of trace atmospheric impurities is obtained from the facts that: (1) ion clusters are readily formed in moist air, (2) the presence of trace impurities tends to form larger, more stable clusters, rapidly, and (3) the smaller diffusivities of the large ionic clusters provide a means of separating the clusters according to size.

Although the IDS responds (with varying sensitivity) to many compounds it is essentially nonresponsive to "normal" atmospheric pollutants (e.g.,  $NO_x$ ,  $SO_2$ ,  $H_2S$ , etc.) and other compounds of low molecular weight. Therefore, interferences from these substances and others likely to be found in a realistic environment of its intended use are minimal. Whatever inadequacies such a system suffers in its lack of specificity for detection are made up in its inherent sensitivity and its simplicity.

### LITERATURE CITED

1. Tickner, A. W., and Knewstubb, P. F. Mass Spectrometric Studies of Ions in Glow Discharges. IV. Water Vapour. *J. Chem. Phys.* **38**, 464 (1963).
2. Kebarle, P., and Godbole, E. W. Mass Spectrometric Study of Gases at Near-Atmospheric Pressures. *J. Chem. Phys.* **39**, 1131 (1963).
3. Narcisi, R. S., and Bailey, A. D. Mass Spectrometric Measurements of Positive Ions at Altitudes from 64 to 112 Kilometers. *J. Geophys. Res.* **70**, 3787 (1965).
4. Kebarle, P. High-Order Reaction - Ion Clusters and Ion Solvation (with references), in *Ion Molecule Reactions*, Edited by J. L. Franklin. Vol. 1, Chap. 7. Plenum Press. New York, New York. 1972.
5. Ineson, T. C., and Harden, C. S. EATR 4642. Chemical Ionization Ion Cluster Mass Spectrometry. May 1972.
6. Carroll, D., *et al.* Subpicogram Detection System for Gas Phase Analysis Based upon Atmospheric Pressure Ionization (API) Mass Spectrometry. *Anal. Chem.* **46**, 706-710 (1974).
7. Good, A., Durden, D. A., and Kebarle, P. Ion Molecule Reactions in Pure Nitrogen and Nitrogen Containing Traces of Water at Total Pressures of 0.5-4 Torr. Kinetics of Clustering Reactions Forming  $H^+(H_2O)_n$ . *J. Chem. Phys.* **52**, 212 (1970).
8. *Ibid.* Mechanism and Rate Constants of Ion-Molecule Reactions Leading to Formation of  $H^+(H_2O)_n$  in Moist Oxygen Air. 222.
9. Ferguson, F. F., and Fehsenfeld, F. C. Water Vapor Ion Cluster Formation in the D Region. *J. Geophys. Res.* **74**, 5743 (1969).
10. Shahin, M. M. Use of Corona Discharges for the Study of Ion-Molecule Reactions. *J. Chem. Phys.* **47**, 4392 (1967).
11. Harden, C. S. EATR 4569. A Mechanism for the Formation of Electrically Charged Ammonia-Water Clusters in the Condensation Nuclei Personnel Detector. November 1971.
12. Anderson, W. F., Schluter, B. C., and Harris, W. A. Honeywell, Inc. Final Report. Contract AFATL-TR-72-212. Multi-Agent Detector System (U). November 1972.

13. Kosolapoff, G. M. Dipole Moments of the Lower Dialkyl Alkylphosphonates. J. Chem. Soc. (London), 3222 (1954).

14. Nelson, R. D., Lide, D. R., and Maryott, A. A. NSARDS-NBS 10. Selected Values of Electric Dipole Moments for Molecules in the Gase Phase. US Government Printing Office, Washington, D.C. 1967.

15. Burke, R. R., and Miller, W. J. AeroChem Research Laboratories, Inc. Third Quarterly Progress Report. Contract DAAA15-68-C-0469. Research Studies on the Condensation Nuclei (CN) Personnel Detector. April 1969.

16. McDaniel, E. W., and Mason, E. A. The Mobility and Diffusion of Ions in Gases. John Wiley and Sons, Inc. New York, New York. 1973.

# DISTRIBUTION LIST 4

Names	Copies	Names	Copies
CHEMICAL SYSTEMS LABORATORY		Defense Intelligence Agency	
SAFETY OFFICE		Attn: DB-4G1	1
Attn: DRDAR-CLF	1	Washington, DC 20301	
AUTHOR'S COPIES: Research Division	10	DEPARTMENT OF THE ARMY	
BIOMEDICAL LABORATORY		HQDA (DAMO-SSC)	1
Attn: DRDAR-CLL-B	1	WASH DC 20310	
Attn: DRDAR-CLL-M	1	Director	
Attn: DRDAR-CLL-MC	1	Defense Civil Preparedness Agency	
Attn: DRDAR-CLL-V	1	Attn: PO(DC)	1
CB DETECTION & ALARMS DIVISION		Washington, DC 20301	
Attn: DRDAR-CLC-C	1	CINCUSAREUR	
Attn: DRDAR-CLC-E	1	Attn: AEAGC-RSI	1
DEVELOPMENTAL SUPPORT DIVISION		APO New York 09403	
Attn: DRDAR-CLJ-R	3	Deputy Chief of Staff for: Research,	
Attn: DRDAR-CLJ-L	3	Development & Acquisition	
Attn: DRDAR-CLJ-M	1	Attn: DAMA-CSM-CM	1
Attn: DRDAR-CLJ-P	1	Attn: DAMA-ARZ-D	1
ENVIRONMENTAL TECHNOLOGY DIVISION		Washington, DC 20310	
Attn: DRDAR-CLT-D	4	US Army Research and Standardization	
MUNITIONS DIVISION		Group (Europe)	1
Attn: DRDAR-CLN	2	Attn: Chief, Chemistry Branch	
PHYSICAL PROTECTION DIVISION		Box 65, FPO New York 09510	
Attn: DRDAR-CLW	2	HQDA (DAMI-FIT)	1
Attn: DRDAR-CLW-C	1	WASH, DC 20310	
Attn: DRDAR-CLW-P	1	Commander	
Attn: DRDAR-CLW-E	1	HQ US Army Medical Command, Europe	
RESEARCH DIVISION		Attn: AEMPM	1
Attn: DRDAR-CLB	1	APO New York 09403	
Attn: DRDAR-CLB-B	1	US ARMY HEALTH SERVICE COMMAND	
Attn: DRDAR-CLB-C	1	Superintendent	
Attn: DRDAR-CLB-P	1	Academy of Health Sciences	
Attn: DRDAR-CLB-R	1	US Army	
Attn: DRDAR-CLB-T	1	Attn: HSA-CIX	1
Attn: DRDAR-CLB-TE	1	Attn: HSA-JHE	1
SYSTEMS DEVELOPMENT DIVISION		Fort Sam Houston, TX 78234	
Attn: DRDAR-CLY-A	1	US ARMY MATERIEL DEVELOPMENT AND	
Attn: DRDAR-CLY-L	4	READINESS COMMAND	
Attn: DRDAR-CLY-R	6	Commander	
Attn: DRDAR-CLY-T	1	US Army Materiel Development and Readiness Command	
DEPARTMENT OF DEFENSE		Attn: DRCLIX	1
Administrator		Attn: DRCSE-P	1
Defense Documentation Center		5001 Eisenhower Ave	
Attn: Document Processing Division (DDC-DD)	12	Alexandria, VA 22333	
Cameron Station			
Alexandria, VA 22314			

# DISTRIBUTION LIST 4 (Contd)

Names	Copies	Names	Copies
Commander US Army Toxic & Hazardous Materials Agency Attn: DRXTH-TS Aberdeen Proving Ground, MD 21010	2	Commander USA Technical Detachment US Naval EOD Facility Indian Head, MD 20640	1
Human Engineering Laboratory HFE Detachment Attn: DRXHE-EA Building E3220 Aberdeen Proving Ground, MD 21010	1	US ARMY ARMAMENT MATERIEL READINESS COMMAND	
Commander US Army Foreign Science & Technology Center Attn: DRXST-MT2 220 Seventh St., NE Charlottesville, VA 22901	1	Commander US Army Armament Materiel Readiness Command Attn: DRSAR-ASN Attn: DRSAR-IRB Attn: DRSAR-PDM Attn: DRSAR-PE Attn: DRSAR-SF Attn: DRSAR-SR Rock Island, IL 61299	1 1 1 1 1 1
US Army Materiel Systems Analysis Activity Attn: DRXSY-MP Attn: DRXSY-T (Mr. Kaste) Aberdeen Proving Ground, MD 21005	1 1	CDR, APG USA ARRCOM Attn: DRSAR-MAS-C Attn: SARTE Aberdeen Proving Ground, MD 21010	 1 1
Commander US Army Missile Research and Development Command Redstone Scientific Information Center Attn: DRDMI-TBD Redstone Arsenal, AL 35809	1	Commander US Army Dugway Proving Ground Attn: Technical Library, Docu Sect Dugway, UT 84022	1
Director DARCOM Field Safety Activity Attn: DRXOS-C Charlestown, IN 47111	1	Commander Rocky Mountain Arsenal Attn: SARRM-QA Attn: SARRM-MD Commerce City, CO 80022	1 1
US ARMY ARMAMENT RESEARCH AND DEVELOPMENT COMMAND		Commander Pine Bluff Arsenal Attn: SARPB-ETA Pine Bluff, AR 71611	1
Commander US Army Armament Research and Development Command Attn: DRDAR-AC Attn: DRDAR-LCE-M Attn: DRDAR-LCH Attn: DRDAR-LCN Attn: DRDAR-LCU Attn: DRDAR-SER Attn: DRDAR-TSS Attn: DRCPM-CAWS-WP Dover, NJ 07801	1 1 1 1 1 1 2 1	US ARMY TRAINING & DOCTRINE COMMAND	
Director Ballistic Research Laboratory Attn: DRDAR-TSD-S Building 305 Aberdeen Proving Ground, MD 21005	1	Commandant US Army Infantry School Attn: NBC Division Fort Benning, GA 31905	1
CDR, APG USA ARRAIDCOM Attn: DRDAR-GCL Aberdeen Proving Ground, MD 21010	1	Commandant US Army Missile & Munitions Center & School Attn: ATSK-CD-MD Attn: ATSK-DT-MU-EOD Redstone Arsenal, AL 35809	1 1
		Commandant US Army Military Police School/Training Center Attn: ATZN-CDM Attn: ATZN-TDPC Fort McClellan, AL 36205	1 1



# DISTRIBUTION LIST 4 (Contd)

Names	Copies	Names	Copies
Commander US Army Infantry Center Attn: ATSH-CD-MS-C Fort Benning, GA 31905	1	US MARINE CORPS Director, Development Center Marine Corps Development & Education Command Attn: Fire Power Division Quantico, VA 22134	1
Commandant US Army Ordnance & Chemical Center & School Attn: ATSL-CL-CD Aberdeen Proving Ground, MD 21005	1	DEPARTMENT OF THE AIR FORCE Air University Library Attn: AUL/LSE-8879 Maxwell AFB, AL 36112	1
US ARMY TEST & EVALUATION COMMAND Commander US Army Test & Evaluation Command Attn: DRSTE-FA Aberdeen Proving Ground, MD 21005	1	HQ Foreign Technology Division (AFSC) Attn: PDRR Wright-Patterson AFB, OH 45433	1
President US Army Infantry Board Attn: ATZB-IB-MI Fort Benning, GA 31905	1	Commander Aeronautical Systems Division Attn: ASD/AELD Wright-Patterson AFB, OH 45433	1
DEPARTMENT OF THE NAVY Chief of Naval Research Attn: Code 443 800 N. Quincy Street Arlington, VA 22217	1	HQ, USAF/SGPR Forrestal Bldg WASH DC 20314	1
Commander Naval Facilities Engineering Command Attn: Code 03 200 Stovall Street Alexandria, VA 22332	1	HQ AFSC/SDAE Andrews AFB, MD 20334	1
Commander Naval Explosive Ordnance Disposal Facility Attn: Army Chemical Officer, Code 604 Indian Head, MD 20640	1	HQ AFISC/SEV Norton AFB, CA 92409	1
Commander Nuclear Weapons Training Group, Atlantic Naval Air Station Attn: Code 21 Norfolk, VA 23511	1	NORAD Combat Operations Center/DBN Cheyenne Mtn Complex, CO 80914	1
Chief, Bureau of Medicine & Surgery Department of the Navy Washington, DC 20372	1	AMREI/MEB Wright-Patterson AFB, OH 45433	1
Commander Naval Weapons Center Attn: Code 3171 China Lake, CA 93555	1	OUTSIDE AGENCIES Battelle, Columbus Laboratories Attn: TACTEC 505 King Avenue Columbus, OH 43201	1
		Director of Toxicology National Research Council 2101 Constitution Ave, NW Washington, DC 20418	1
		ADDITIONAL ADDRESSEES Commander USEUCOM Attn: ECJS-O/LTC James H. Alley APO, NY 09128	1
		Commander Armament Concepts Office, Weapon Systems Concepts Team Attn: DRDAR-ACW Aberdeen Proving Ground, MD 21010	1